arXiv:1209.4036v3 [quant-ph] 20 Sep 2012

Novel States of Classical Light and Noncontextuality

Partha Ghose*
Centre for Astroparticle Physics and Space Science (CAPSS),
Bose Institute,
Block EN, Sector V, Salt Lake, Kolkata 700 091, India,
and
Anirban Mukherjee,
Indian Institute of Science Education & Research,
Mohanpur Campus, West Bengal 741252.

PACS nos. 42.25.-p, 42.25.Ja, 42.25.Kb, 03.65.Ud

Keywords: classical light, polarization, Pancharatnam phase, entanglement, nonseparability, noncontextuality

Abstract

Novel states of classical light that are nonseparable are introduced, and experiments are proposed to test the violation of noncontextuality by these Bell-like states. Such states have only recently begun to be explored. The significance of their nonseparability or entanglement as well as their similarities with and differences from entangled quantum states are analysed. A separate experiment is proposed to verify the predicted variation of nonseparability of classical light with the Pancharatnam phase.

---

*partha.ghose@gmail.com

1 Introduction

Entanglement has been held to be *the* trait of quantum mechanics that characterises its entire departure from classical physics [1]. If non-factorizability of independent states of a system is taken as the defining characteristic of their loss of identity through entanglement, then entanglement of classical light and the consequent violation of Bell-like inequalities, without implying nonlocality, are inevitable consequences of its Hilbert space structure, as was shown way back in 2001 [2]. In fact, birefringent crystals have been known since the 17th century to produce states of the form $Au\otimes s+Bv\otimes p$ where u is the path with s polarization and v with p polarization, A and B being amplitudes. Recently, the essential role of non-quantum entanglement in resolving a certain basic issue in classical polarization optics has been pointed out [3], and nonseparable cylindrically polarized laser beams have been extensively studied and used [4, 5, 6]. The predicted violation of a Bell-like inequality by polarization-path nonseparable/entangled classical states of light has also been experimentally verified [7, 8].

There is no doubt that there are some crucial differences between entanglement/nonseparability in classical optics and the more familiar entanglement in quantum mechanics. Hence, in order not to confuse the two, various suggestions have been made about the correct nomenclature to be used in the case of classical light. Some have suggested the use of 'nonseparability' [7], some 'structural inseparability' [8] and some 'non-quantum entanglement' [3]. We would prefer to use 'classical entanglement' to distinguish it from 'quantum entanglement' but will also use nonseparability equivalently. The reason is to emphasize the fact that non-factorizability already occurs in classical electrodynamics because of its Hilbert space structure which is not unique to quantum mechanics. Hence, classical electrodynamics is more like quantum mechanics than is Newtonian mechanics, and a classical-quantum transition is likely to be better understood if classical electrodynamics is taken as the fiducial classical theory. One can then go directly from relativistic classical field theory to relativistic quantum field theory and vice versa, and then take appropriate limits to get relativistic single-particle quantum mechanics and finally non-relativistic quantum mechanics. The choice of the other path, namely from Newtonian mechanics to relativistic quantum field theory, has been plagued by many problems like entanglement that refuse to go away.

A big advantage of classical entanglement is that it is robust whereas quantum entanglement is notoriously fragile. This fact also needs to be better understood. From the practical point of view alone, many manipulations that are necessary for quantum information processing (but not all) can be done far more easily with classical light than with quantum light.

Another very important aspect of classical entanglement is its bearing on the concept of noncontextuality and realism. This has not been discussed at all so far. In this paper we wish to go into this question and show that the polarization and spatial modes of classically entangled light are *contextual* variables, and have no value definiteness, which is a real surprise in classical physics.

In Section 2, a simple way to produce all four Bell-like states of polarization-path entangled classical light will be described, and the state $|\Phi^+\rangle$ will be shown to violate a Bell-like inequality derived from noncontextuality [9]. Violation of the axioms of the Kochen-Specker theorem [10, 11, 12] by the state $|\Phi^-\rangle$ will also be shown. Classically entangled two-mode states analogous to two-particle quantum states will be introduced and their implications discussed in Section 3. The variation of nonseparability of a Bell-like classical state with the Pancharatnam phase will be studied in Section 4, and an experiment proposed to test this prediction. The important differences between classical and quantum entanglement will be discussed in Section 5.

2 Bell-like States in Classical Optics and Noncontextuality

Classical electrodynamics is the paradigm of classical field theories in physics. That classical electrodynamics has a Hilbert space structure was first explicitly shown in 2001 [2]. Without getting involved in the details of that demonstration, let us simply note that two different Hilbert spaces are required for a complete description of an ordinary state in classical electrodynamics, namely a space $\hat{H}_{path}$ of square integrable functions that describe scalar optics and a two-dimensional space of polarization states $\hat{H}_{pol}$. These are disjoint Hilbert spaces, and hence a complete description of a state is given in terms of tensor products of states in these two Hilbert spaces: $\frac{1}{\sqrt{|A|^2}}|A\rangle \otimes |\lambda\rangle \in \hat{H}_{path} \otimes \hat{H}_{pol}$ where $A(\mathbf{r}, t) = \langle \mathbf{r}, t|A\rangle$ are solutions of the scalar wave equation

$$\left[\nabla^2 - \frac{1}{c^2}\frac{\partial^2}{\partial t^2}\right] A(\mathbf{r}, t) = 0 \tag{1}$$

and $|\lambda\rangle$ is the vector

$$|\lambda\rangle = \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix}$$

of the transverse polarizations λ_1 and λ_2. $A(\mathbf{r}, t)$ is a Laguerre-Gauss polynomial for laser beams. One can also write $\frac{1}{\sqrt{|A|^2}}|A\rangle \otimes |\lambda\rangle$ more conventionally as the Jones vector

$$|J\rangle = \frac{1}{\sqrt{\langle J|J\rangle}} \begin{pmatrix} E_x \\ E_y \end{pmatrix}$$

where $E_x = A_0 \hat{e}_x \exp(\mathrm{i}\phi_\mathrm{x})$ and $E_y = A_0 \hat{e}_y \exp(\mathrm{i}\phi_\mathrm{y})$ are the complex transverse electric fields, $\hat{e}_x$ and $\hat{e}_y$ are unit polarization vectors, and $\langle J|J\rangle = |E_x|^2 + |E_y|^2 = A_0^2$ is the intensity I_0. Given this mathematical structure of a tensor product Hilbert space, polarization-path entanglement is inevitable, and one can construct the complete set of Bell-like states.

Let us consider a normalized state $|\psi\rangle_{in} = (A/\sqrt{I_0})|a_{in}\rangle \otimes |V\rangle$ of an incoming vertically polarized monochromatic classical light beam of intensity $I_0 = |A|^2$. An example would be a laser beam that does not show any quantum effect like squeezing or sub-Poissonian statistics, and of coherence length longer than the paths through the interferometer. When such a light beam is incident on a 50-50 lossless non-polarizing beam splitter $NPBS_1$ (Fig. 1), the transmitted beam $|a\rangle$ and the reflected beam $|b\rangle$ span a two-dimensional Hilbert space $\hat{H}_{path} = \{|a\rangle, |b\rangle\}$. Another two-dimensional Hilbert space $\hat{H}_{pol} = \{|V\rangle, |H\rangle\}$ is associated with the polarization states of the two ougoing light beams. Hence the full Hilbert space is $\hat{H} = \hat{H}_{path} \otimes \hat{H}_{pol}$. The action of $NPBS_1$ on the incident beam can be described by the unitary matrix

$$U^{path}_{NPBS} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & i \\ i & 1 \end{pmatrix}_{path} \otimes \mathbb{I}^{pol} \tag{2}$$

where $\mathbb{I}^{pol}$ is the unit operator on $\hat{H}_{pol}$, resulting in the state

$$U^{path}_{NPBS}|\psi\rangle_{in} = \frac{A}{\sqrt{2I_0}} \left[|a\rangle \otimes |V\rangle + i|b\rangle \otimes |V\rangle \right]. \tag{3}$$

When this is followed by a quarter-wave phase-shifter plate in path b with its fast axis vertical, represented by the Jones matrix

$$J_{QWP} = \begin{pmatrix} 1 & 0 \\ 0 & -i \end{pmatrix} \tag{4}$$

which acts on $\hat{H}_{pol}$, the unitary matrix U that acts on the state is

$$U = |a\rangle\langle a| \otimes \mathbb{I}^{pol} + |b\rangle\langle b| \otimes \hat{J}_{QWP}, \tag{5}$$

resulting in the output state

$$\begin{aligned} |\Psi_{out1}\rangle &= \frac{A}{\sqrt{2I_0}}\left[|a\rangle \otimes |V\rangle + i|b\rangle \otimes J_{QWP}|V\rangle\right] \\ &= \frac{A}{\sqrt{2I_0}}\left[|a\rangle \otimes |V\rangle + |b\rangle \otimes |V\rangle\right]. \end{aligned} \tag{6}$$

The action of the polarization flipper PF_b on this state can be represented by the unitary matrix

$$U_{flip} = |a\rangle\langle a| \otimes \mathbb{I}^{pol} + |b\rangle\langle b| \otimes \hat{J}_{flip} \tag{7}$$

where $\hat{J}_{flip}$ is the Jones matrix with $J_{xx} = J_{yy} = 0, J_{xy} = J_{yx} = 1$, so that $\hat{J}|V\rangle = |H\rangle$, and the final state produced is

$$|\Phi^+\rangle = U_{flip}|\Psi_{out1}\rangle = \frac{A}{\sqrt{2I_0}}\left[|a\rangle \otimes |V\rangle + |b\rangle \otimes |H\rangle\right]. \tag{8}$$

This is a polarization-path entangled Bell-like state. One can introduce a π phase shifter in the path b to get the state $|\Phi^-\rangle$. Similarly, one can get the other two Bell states $|\Psi^+\rangle$ and $|\Psi^-\rangle$ from a beam $|A\rangle \otimes |H\rangle$.

Our next aim is to test whether these states are consistent with noncontextuality which is a straightforward generalization of an apparently innocuous realism about measurement, namely the conventional classical notion that a physical reality must be independent of how it is measured. Put more technically, noncontextuality is the requirement that the result of a measurement is predetermined and not affected by how the value is measured, i.e. not affected by previous or simultaneous measurement of any other compatible or co-measureable observable. To test that, joint measurements of compatible observables that are not necessarily spatially separated are required. This can be done for classical light by making a joint measurement of its two independent degrees of freedom, namely path and polarization, in one beam path. If the light is prepared in a non-factorizable state like $|\Phi^+\rangle$, one can compare the predicted correlations with a Bell-like inequality for such correlations derived from the requirement of noncontextuality alone [9] without any reference to Einstein locality. This is what we proceed to do now.



Figure 1: Schematic diagram of the experiment: A vertically polarized beam of classical light passes through a Mach-Zehnder interferometer. PF_b in path b is a polarization flipper that converts V to H. PS_b is a phase shifter and PR_b is a polarization rotator in path b.

Let M_a and M_b be two lossless mirrors (Fig. 1) and $NPBS_2$ a second 50-50 lossless nonpolarizing beam splitter which is followed by a single light detector in the path a_{out}. Also, let a phase shifter PS_b and a polarization rotator PR_b be inserted in path b as shown in Fig. 1. The polarization rotator matrix is $R(\theta) = \exp(i\theta)\hat{J}_{QWP}$, i.e. a variable phase-shifter combined with a quarter-wave plate with its fast axis vertical or horizontal. Then, the state before $NPBS_2$ is

$$\frac{iA}{\sqrt{2I_0}}[|a\rangle \otimes |V\rangle + e^{i\phi}|b\rangle \otimes e^{i\theta}|H\rangle]. \tag{9}$$

The final state after $NPBS_2$ is

$$\begin{aligned} |\Phi^+\rangle' &= \frac{iA}{2\sqrt{I_0}}[|b_{\text{out}}\rangle \otimes (|V\rangle + ie^{i(\theta+\phi)}|H\rangle) \\ &+ |a_{\text{out}}\rangle \otimes (i|V\rangle + e^{i(\theta+\phi)}|H\rangle)]. \end{aligned} \tag{10}$$

Now, consider the projection operators

$$\sigma_\theta = \mathbb{I}^{path} \otimes \left(2|\text{pol}\rangle\langle\text{pol}| - \mathbb{I}^{pol}\right), \tag{11}$$

$$\sigma_\phi = \left(2|\text{path}\rangle\langle\text{path}| - \mathbb{I}^{path}\right) \otimes \mathbb{I}^{pol}, \tag{12}$$

with

$$
\begin{aligned}
|\text{path}\rangle &= \frac{1}{\sqrt{2}}[|a\rangle + e^{i\phi}|b\rangle], \qquad (13)\\
|\text{pol}\rangle &= \frac{1}{\sqrt{2}}[|V\rangle + e^{i\theta}|H\rangle]. \qquad (14)
\end{aligned}
$$

Their eigenvalues are ±1. Hence,

$$
\begin{aligned}
E(\theta,\chi) &= \langle\Phi^+|\sigma_\theta\,.\,\sigma_\phi|\Phi^+\rangle\\
&= \langle\Phi^+|\,[(+)\sigma_{\theta,+1} + (-)\sigma_{\theta,-1}].[(+)\sigma_{\phi,+1} + (-)\sigma_{\phi,-1}]|\Phi^+\rangle\\
&= \cos(\theta+\phi) \qquad (15)
\end{aligned}
$$

where

$$
\begin{aligned}
\sigma_{\theta,\pm 1} &= (|V\rangle \pm e^{i\theta}|H\rangle)(\langle V| \pm e^{-i\theta}\langle H|) \qquad (16)\\
\sigma_{\phi,\pm 1} &= (|a\rangle \pm e^{i\phi}|b\rangle)(\langle a| \pm e^{-i\phi}\langle b|). \qquad (17)
\end{aligned}
$$

$E(\theta,\phi)$ is thus the result of compatible joint measurements of the states $|\text{pol}\rangle$ and $|\text{path}\rangle$.

Now define

$$
S = E(\theta_1,\phi_1) + E(\theta_1,\phi_2) - E(\theta_2,\phi_1) + E(\theta_2,\phi_2). \qquad (18)
$$

By requiring that the polarizations and paths have predetermined values in a classical state that do not depend on whether or not one of them is being measured or both, simultaneously or at different times, a Bell-like inequality can be expressed as $-2 \leq S \leq 2$ [9]. This is obviously violated by the state $|\Phi^+\rangle$ for the set $\theta_1 = 0, \theta_2 = \pi/2, \phi_1 = \pi/4, \phi_2 = -\pi/4$ for which $S = 2\sqrt{2}$.

Since the path and polarization changes are made on the same state in path b, there is no question of any nonlocality in this experiment. The predicted violation therefore shows that the path and polarization of even classical light in Bell-like states must be contextual. This is analogous to the case of path-spin polarization in neutron optics [15].

One can also show that these classical Bell-like nonseparable states violate the axioms of noncontextual hidden variable theories on which the Kochen-Specker theorem is based [10, 11, 12]. Consider the state

$$
|\Phi^-\rangle = \frac{A}{\sqrt{2I_0}}\,[|a\rangle \otimes |V\rangle - |b\rangle \otimes |H\rangle] \qquad (19)
$$

and six hermitian operators or 'observables' $J_x^{pol}, J_x^p, J_y^{pol}, J_y^p, J_x^{pol}J_y^p, J_y^{pol}J_x^p$ acting on it, where J^{pol}s are Jones matrices acting on $\hat{H}_{pol}$ and J^ps are analogous matrices acting on $\hat{H}_{path}$. Hence, J^{pol}s and J^ps commute. These matrices are taken to have the same form as the Pauli matrices σ_i with the property $\sigma_x\sigma_y = i\sigma_z$. Hence, the six operators have eigenvalues ±1. Since

$$\begin{aligned} J_x^p|a\rangle &= |b\rangle, \quad J_x^p|b\rangle = |a\rangle, && (20)\\ J_y^p|a\rangle &= i|b\rangle, \quad J_y^p|b\rangle = -i|a\rangle, && (21)\\ J_x^{pol}|V\rangle &= |H\rangle, \quad J_x^{pol}|H\rangle = |V\rangle, && (22)\\ J_y^{pol}|V\rangle &= -i|H\rangle, \quad J_y^{pol}|H\rangle = i|V\rangle, && (23) \end{aligned}$$

it follows that

$$\begin{aligned} J_x^{pol} \,.\, J_x^p|\Phi^-\rangle &= -|\Phi^-\rangle, && (24)\\ J_y^{pol} \,.\, J_y^p|\Phi^-\rangle &= -|\Phi^-\rangle, && (25)\\ J_x^{pol}J_y^p \,.\, J_x^{pol} \,.\, J_y^p|\Phi^-\rangle &= +|\Phi^-\rangle, && (26)\\ J_y^{pol}J_x^p \,.\, J_y^{pol} \,.\, J_x^p|\Phi^-\rangle &= +|\Phi^-\rangle, && (27)\\ J_x^{pol}J_y^p \,.\, J_y^{pol}J_x^p|\Phi^-\rangle &= -|\Phi^-\rangle. && (28) \end{aligned}$$

Since each of the six observables occurs exactly twice on the left-hand sides, the product of the left-hand sides is $+1$ on the assumption that each observable has a predetermined value ±1, whereas the product of the right-hand sides is -1. This is a clear logical contradiction. The Kochen-Specker theorem is based on two assumptions, namely (i) value definiteness, i.e. all observables of a system have predetermined values, and (ii) noncontextuality which requires these values to be independent of the way in which they are measured. Hence, these assumptions cannot hold for nonseparable classical states like $|\Phi^-\rangle$. An experiment can be designed to test this prediction along the lines of the neutron experiment of Hasegawa et al [13].

3 Nonseparable Two-mode Classical States

Consider a laser beam in the double mode $|H\rangle^a \otimes |H\rangle^b$ where each ket represents a first-order Hermite-Gaussian transverse mode $\psi_H(\vec{r})^{a/b}\hat{e}_H$ with horizontal orientation in a path (a or b), $\hat{e}_H$ being the horizontal (H) linear polarization unit vector. Similarly, let $|V\rangle^c \otimes |V\rangle^d$ be a second laser double



Figure 2: Combining two double-mode beams, one in the VV and the other in the HH mode.

mode with vertical orientation in paths c and d. They are product states. Particular cases for $\psi_H(\vec{r})$ and $\psi_V(\vec{r})$ would be the modes $\mathrm{TEM}_{01(y)}$ and $\mathrm{TEM}_{01(x)}$ respectively. It is possible to combine these beams and produce nonseparable linear combinations as shown in Figure 2. Let us assume that the four modes have a common intensity $I_0 = |A|^2$. Consider one such state

$$|\Phi^+\rangle = \frac{A}{\sqrt{2I_0}} \left[|H\rangle^a \otimes |H\rangle^b + |V\rangle^a \otimes |V\rangle^b \right]. \tag{29}$$

It is an analog of the two-particle Bell state $|\Phi^+\rangle$ in quantum mechanics. Clearly, all four Bell-like states can be produced in this way. To see what it implies for the theory, let us assume that the modes travel in paths a and b in opposite directions along the z axis, and let us write the state (29) cryptically in the form

$$|\Phi^+\rangle = \frac{1}{\sqrt{2}}\left[\hat{x}^a\hat{x}^b + \hat{y}^a\hat{y}^b\right]. \tag{30}$$

Let a polarization analyzer be placed in each path such that their polarization axes make arbitrary angles α and β with the x axis, and let $\alpha - \beta = \theta$. Let us define the projection operator for linear polarization by

$$P_\theta = (x\cos\theta + y\sin\theta)(x\cos\theta + y\sin\theta)^\dagger, \tag{31}$$

and from it construct the projector

$$\sigma_\theta = 2P_\theta - \mathbb{I} = (\hat{x}\hat{x}^\dagger - \hat{y}\hat{y}^\dagger)\cos 2\theta + (\hat{x}\hat{y}^\dagger + \hat{y}\hat{x}^\dagger)\sin 2\theta \tag{32}$$

whose eigenvalues are ±1. Then the polarization correlation in the state (29) is given by

$$\langle\sigma_0 \otimes \sigma_\theta\rangle = \cos 2\theta. \tag{33}$$

Therefore, if Alice has the choice of angles α and γ on one side and Bob the choice β and γ on the other side , Bell's inequality, based on local realism, takes the form

$$|\cos 2(\alpha - \beta) - \cos 2(\alpha - \gamma)| + \cos 2(\beta - \gamma) \leq 1 \tag{34}$$

which is violated by the correlation (33) for relative angles of $\pi/6$ between the analyzers, exactly as in quantum mechanics.

Does that mean that such states violate locality? The answer is decidedly no, because there is no collapse of the state on local measurements of polarization, as in standard quantum mechanics, and hence no nonlocal effect. All that happens in a classical measurement is that one of the polarization states is projected (i.e. selected) and recorded while the other is simply blocked– it does not disappear as in projective quantum measurement. Violation of the inequality therefore only shows that, like their counterparts in quantum mechanics, these states are more correlated than states based on classical mechanical notions of local realism.

Multimode lasers can be used in principle to produce multimode nonseparable states of classical light.



Figure 3: The closed circuit on the Poincare sphere

4 Influence of the Pancharatnam Phase on a Classical Bell-like State

When the polarization of a light beam is rotated in a cyclic path, the polarization vector comes back with an overall phase that is the sum of the dynamical phase and the Pancharatnam phase [16] which is half the solid angle subtended by the circuit at the centre of the Poincare sphere. To isolate the Pancharatnam phase, the effect of the dynamical phase has to be isolated, and this can be done in the following manner. Let a half-wave plate (HWP) be inserted between two co-aligned quarter-wave plates (QWPs) to form a combination QHQ. Let such a device be inserted into one arm of an interferometer. It can be shown [17] that if plane polarized light of phase α is incident on the first QWP at 45^0 to its axes, its phase shifts to $\alpha + \pi - 2\gamma$ where π is the dynamical phase shift due to the optical path lengths and γ is the angle between the axes of the HWP and the QHQ. As the HWP is rotated, γ changes, and this change is detectable through fringe shifts of -2γ. This is a pure geometrical phase shift brought about by polarization changes, and is hence the Pancharatnam phase. The closed circuit on the Poincare sphere is shown in Figure 3. Let us now consider the experimental arrangement shown in Figure 4 which is a modification of Fig. 1. Let the first beam splitter $NPBS_1$ be replaced by a variable asymmetric nonpolariz-



Figure 4: Fig. 1 modified by replacing $NPBS_1$ by a variable $VNPBS_1$ and adding optical elements QHQ and HWP_a in path a and a FWP_b and HWP_b in path b.

ing beam splitter $VNPBS_1$ with the reflection and transmission coefficients $r = \sin\theta_1$ and $t = \cos\theta_1$, and let a QHQ and a half-wave plate HWP_a be inserted in path a and a full-wave plate FWP_b and a half-wave plate HWP_b be inserted in path b. Fixing the initial arrangement so that $\theta_1 = \pi/4$, the state after QHQ in path a and FWP_b in path b s

$$|\Phi_P\rangle = \frac{Ae^{i\pi}}{\sqrt{2I_0}}[e^{-2i\gamma}|a\rangle|V\rangle + |b\rangle|H\rangle]. \tag{35}$$

Then, after passage through the other optical elements but before the second beam splitter $NPBS_2$ and for variable θ_1, the state is given by

$$\begin{aligned} |\Phi\rangle &= \frac{Ae^{i\pi}}{\sqrt{I_0}}[e^{-i2\gamma}\cos\theta_1|a\rangle \otimes (\sin\theta_2|H\rangle - \cos\theta_2|V\rangle) \\ &+ e^{i\phi}\sin\theta_1|b\rangle \otimes \big(\cos\theta_2|H\rangle + e^{i\theta}\sin\theta_2|V\rangle\big)] \end{aligned} \tag{36}$$

where HWP_a and HWP_b have their fast axes at $\theta_2/2$ to the vertical. After passing $NPBS_2$, the final state is

$$\begin{array}{rcl}|\Phi\rangle & = & \frac{Ae^{i\pi}}{\sqrt{I_0}}[|a\rangle \otimes \big\{\cos\theta_1 \sin\theta_2 + ie^{i\phi}\sin\theta_1\cos\theta_2\big)\,|H\rangle \\ & + & \big(ie^{i(\theta+\phi)}\sin\theta_2\sin\theta_1 - e^{-2i\gamma}\cos\theta_1\cos\theta_2\big\}\,|V\rangle \\ & + & |b\rangle \otimes \big\{\big(ie^{-2i\gamma}\cos\theta_1\sin\theta_2 + e^{i\phi}\sin\theta_1\cos\theta_2\big)\,|H\rangle\big) \\ & + & \big(e^{i(\theta+\phi)}\sin\theta_2\sin\theta_1 - ie^{-2i\gamma}\cos\theta_1\cos\theta_2\big)\,|V\rangle\}]. \end{array} \tag{37}$$

Therefore, using the projection operators (11) and (12) with

$$|\text{path}\rangle = \cos\theta_1|a\rangle + e^{i\phi}\sin\theta_1|b\rangle, \tag{38}$$

$$|\text{pol}\rangle = \cos\theta_2|V\rangle + e^{i\theta}\sin\theta_2|H\rangle, \tag{39}$$

we get

$$E\left(\theta_1,\theta_2,\theta,\phi,\gamma\right) = \langle\sigma_\theta\,.\,\sigma_\phi\rangle = \cos 2\theta_1\cos 2\theta_2 + \cos(\theta+\phi-2\gamma)\sin 2\theta_1\sin 2\theta_2. \tag{40}$$

One can then define the function

$$\begin{array}{rcl} S & = & E(\theta_1,\theta_2,\theta,\phi,\gamma) - E(\theta_1^{'},\theta_2,\theta,\phi^{'},\gamma) + E(\theta_1,\theta_2^{'},\theta^{'},\phi,\gamma) \\ & + & E(\theta_1^{'},\theta_2^{'},\theta^{'},\phi^{'},\gamma) \end{array} \tag{41}$$

which must satisfy the Bell-like inequality $-2 \leq S \leq 2$. Putting $\theta_2 = 0$ and $\theta = 0$ we have,

$$\begin{array}{rcl} S & = & \cos\left(2\theta_1\right) - \cos 2\theta_1' + \cos\left(2\theta_1\right)\cos\left(2\theta_2'\right) + \cos\left(\theta'+\phi-2\gamma\right)\sin\left(2\theta_1\right)\sin\left(2\theta_2'\right) \\ & + & \cos 2\theta_1'\cos\left(2\theta_2'\right) + \cos\left(\theta'+\phi'-2\gamma\right)\sin\left(2\theta_1'\right)\sin\left(2\theta_2'\right). \end{array} \tag{42}$$

Setting the polar bell-angles at $\theta_1' = \frac{3\pi}{8}, \theta_1 = \frac{\pi}{8}, \theta_2' = \frac{\pi}{4}$, one gets

$$S = \sqrt{2} + \frac{1}{\sqrt{2}}\left[\cos\left(\theta'+\phi'-2\gamma\right) + \cos\left(\theta'+\phi-2\gamma\right)\right]. \tag{43}$$

With $\phi' = \phi$ and $\theta' + \phi = 2\gamma$, $S = 2\sqrt{2}$, i.e. there is maximal violation of the Bell-like inequality.

Let us next look at the reverse problem, i.e. calculate the extremum values of S as a function of the Pancharatnam phase γ for fixed azimuthal angles $\{\theta' = \theta = \phi' = \phi = 0\}$. Then,

$$\begin{array}{rcl} S & = & \cos 2\theta_1 - \cos 2\theta_1' + \cos 2\theta_2'\left(\cos 2\theta_1 + \cos 2\theta_1'\right) \\ & + & \cos 2\gamma\sin 2\theta_2'\left(\sin 2\theta_1 + \sin 2\theta_1'\right). \end{array} \tag{44}$$



h

Figure 5: The polar angle θ_1 is plotted against the Pancharatnam phase γ



h

Figure 6: The polar angle θ_1' is plotted against the Pancharatnam phase γ



h

Figure 7: The extrema of the S function as a function of the Pancharatnam phase for fixed azimuthal angles $\{\theta' = \theta = \phi' = \phi = 0\}$

By requiring the partial derivatives of S w.r.t $\theta_1, \theta_1', \theta_2'$, to vanish,

$$\begin{aligned}
\frac{\partial S}{\partial \theta_1} &= -2\sin 2\theta_1 - 2\sin 2\theta_1 \cos 2\theta_2' + 2\cos 2\gamma \sin 2\theta_2' \cos 2\theta_1 = 0, && (45)\\
\frac{\partial S}{\partial \theta_1'} &= -2\sin 2\theta_1' \cos 2\theta_2' + 2\sin 2\theta_1' + 2\cos 2\theta_1 \cos 2\gamma \sin 2\theta_2' = 0, && (46)\\
\frac{\partial S}{\partial \theta_2'} &= -2\sin 2\theta_2' (\cos 2\theta_1 + \cos 2\theta_1') \\
&+ 2\cos 2\gamma \cos 2\theta_2' (\sin 2\theta_1 + \sin 2\theta_1') = 0, && (47)
\end{aligned}$$

one gets

$$\begin{aligned}
\theta_1 &= \frac{1}{2}\arctan\left(\cos\left(2\gamma\right)\right), && (48)\\
\theta_1^{'} &= \frac{\pi}{2} - \theta_1, && (49)\\
\theta_2' &= \frac{\pi}{4} && (50)
\end{aligned}$$

as the angles for which S is extremal. The results are shown in Figures 5-7. The function S exceeds the Bell limit $+2$ for all values of γ except at the points where it has its minima (Fig. 7).

5 Implications and Significance

Although there is no doubt that quantum entanglement has a much richer structure than classical entanglement, it is clear from the foregoing discussions that that there is more to classical electrodynamics than meets the eye. Classical electrodynamics and quantum mechanics both have a Hilbert space structure [2], which gives rise to many novel features of classical optics that have only recently begun to be explored. It is important to emphasize that there are many examples of the occurrence of Hilbert spaces in classical physics. Euclidean space, for example, is a Hilbert space, the space $\mathbb{C}^n$ of n-tuples of complex numbers $z = (z_1, z_2, ..., z_n)$ and $z' = (z'_1, z'_2, ..., z'_n)$ is a Hilbert space under the inner product defined by $\langle z, z' \rangle = \sum_{k=1}^{n} z_k \bar{z}'_k$. The square integrable functions on L^2 also form a Hilbert space. For any f and g in L^2 one defines the inner product $\langle f, g \rangle = \int_\Omega f(x)\overline{g(x)}dx$. Hence, $\langle f, f \rangle < \infty$. There is nothing 'quantum' about these spaces.

The superposition principle resulting in interference phenomena in classical optics is a straightforward consequence of its Hilbert space structure. There is a mathematical theorem which states that *every pair of vector spaces has a tensor product* [18]. The tensor product space is also a linear vector space. States of classical light are tensor products of two linear vector spaces, the Hilbert space of space-time functions (scalar optics) and the Hilbert space of polarization optics. Hence, *the existence of tensor product spaces resulting in polarization-path entanglement is just as inevitable in classical optics as in quantum mechanics.* It should be clear from this that classical optics (electrodynamics) is a lot more like quantum mechanics than is classical mechanics in which states are points in phase space.

Where, then, are the crucial differences between quantum mechanics and classical optics? The main difference lies in the fact that physical states in quantum mechanics are of unit norm and therefore lie on a unit sphere in Hilbert space. This is necessary for the probabilistic interpretation. Moreover, all states that differ only by an overall phase factor are identified, and hence quantum nechanics actually operates on coset spaces. This is not true of states in classical electrodynamics, resulting in a significant difference between a superposition of states in quantum mechanics and that in classical optics. In quantum mechanics a physical state in general does not *possess* physical properties *before* measurement. For example, a single-photon state like $|X\rangle = c_1|V\rangle + c_2|H\rangle$ with $|c_1|^2 + |c_2|^2 = 1$ cannot be said to possess any polarization V or H. On measurement, however, the state is projected to either $|V\rangle$ with probability $|c_1|^2$ or $|H\rangle$ with probability $|c_2|^2$. This is not the case in classical optics which is deterministic and in which a state $|X\rangle = c_1|V\rangle + c_2|H\rangle$ of light with classical amplitudes c_1 and c_2 *always* possesses a definite polarization given by the vector sum of $c_1|V\rangle$ and $c_2|H\rangle$. $|c_1|$ and $|c_2|$ are here measures of the membership weights of $|X\rangle$ in $|V\rangle$ and $|H\rangle$. This *innocuous scientific realism* of the polarization states in classical optics, however, holds only for product states like $(c_1|x\rangle + c_2|y\rangle) \otimes |w\rangle$ but not for a superposition of product states like $c_1|H\rangle \otimes |x\rangle + c_2|V\rangle \otimes |y\rangle$ in which the polarization and path degrees of freedom are entangled. Such states cannot be said to possess either a definite path or a definite polarization. Hence, these states are of fundamental significance in classical optics, though they have not been considered possible until recently. They encode correlations between polarization states and path states similar to path-spin entangled states in quantum mechanics. They will therefore violate a Bell-like inequality derived from the requirement of noncontextuality, as shown in Section 2 for $|\Phi^+\rangle$. Hence, polarization and path are *contextual* degrees of freedom for such classical light. They do not, of course, imply any nonlocality.

Another fundamental difference between classical and quantum states is that independent quantum states $|A\rangle$ and $|B\rangle$ are decribed by tensor products $|A\rangle \otimes |B\rangle$, but in free space electrodynamics independent light beams are decribed by the sum $|A\rangle \oplus |B\rangle$. Hence, classical states analogous to multiparticle entangled states cannot be produced in free space. They can, however, be produced in laser sources, as we have seen for the case of two-mode beams, and consequently, multi-beam classical entanglement is possible in principle. Again, the difference with quantum entanglement is that the classical states do not violate locality. This raises an important issue of interpretation of empirical data—can the violation of Bell inequalities by quantum states be any longer interpreted as incontrovertible evidence of nonlocality? Local realism is, after all, a special case of noncontextuality where noncommuting observables are measured at space-like intervals to make them mutually compatible.

In spite of the differences, the similarity between quantum mechanics and classical electrodynamics is unexpectedly striking. On the supposition that quantum theory is a more fundamental theory than classical field theory, it is clear that entanglement survives in the classical limit of the theory. The quantum-classical transition is therefore much clearer if field theory is taken as the paradigm of classical physics rather than classical mechanics. *Classical mechanical notions of realism, noncontextuality and separability are as* *impossible to reconcile with quantum mechanics as with classical field theory*. To hold on to realism in some form in classical physics as a whole, one has therefore to search for a deeper and more subtle meaning of reality than is captured in noncontextuality.

Note added: After this paper was uploaded to the arXiv (quant-ph/1209. 4036v2, 19 Sep 2012) Eugenio Roldán pointed out to us the following papers by R. J. C. Spreeuw, namely, *Found. Phys.* **28** (1998) 361 and *Phys. Rev. A* **63** (2001) 062302, which are based on the same basic idea as our paper. However, Spreeuw does not go into the deeper implications of nonseparability or entanglement in classical optics such as noncontextuality and its tests, which are the main concern in this paper.

6 Acknowledgement

We thank C. S. Unnikrishnan, Dipankar Home, Arun Pati, Ujjwal Sen and Tony Sudbery for helpful comments on the preliminary drafts of the paper. The views expressed in this paper are, however, entirely ours. PG thanks the National Academy of Sciences, India for the award of a Senior Scientist Platinum Jubilee Fellowship which allowed this work to be undertaken.

References

[1] Schrödinger, E., *Proc. of the American Phys. Soc.* **124** (1935) 323-38. Reprinted in *Quantum Theory of Measurement*, eds. J. A. Wheeler & W. H. Zurek, Princeton University Press, New Jersey, 1983.

[2] Ghose, P. & Samal, M. K., arXiv:quant-ph/0111119v1 22Nov 2001.

[3] Simon, B. N., Simon, S., Gori, F., Santarsiero, M., Borghi, R., Mukunda, N.& Simon, R., *Phys. Rev. Lett.* **104**, 023901 (2010).

[4] Oron, R., Blit, S., Davidson, N. & Friesem, A. A., *Applied Phys. Lett.* **77**, 3322-3324 (2000).

[5] Tidwell, S. C., Kim, G. H., & Kimura, W. D., *Applied Phys. Lett.* **32**, 5222-5229 (1993).
Kozawa, Y. & Sato, S., *Optics Lett.* **30**, 3063-3065 (2005).

[6] Holleczek, A., Aiello1, A., Gabriel1, C., Marquardt, C., Leuchs, G. (2010). arXiv:1012.4578v1 [quant-ph] 21 Dec 2010.

[7] Borges, C. V. S., Hor-Meyll, M., Huguenin, J. A. O. & Khoury, A. Z., *Phys. Rev. A* **82**, 033833 (2010).

[8] Gabriel, C., Aiello, A., Zhong, W., Euser, T. G., Joly, N. Y., Banzer, P., Fortsch, M., Elser, D., Andersen, U. L., Marquardt, C., Russell, P. S. J.& Leuchs, G., *Phys. Rev. Lett.* **106**, 060502 (2011).

[9] Basu, S., Bandyopadhyay, S., Kar, G., & Home, D., *Phys. Lett. A* **279**, 281-286 (2001).

[10] Kochen, S., Specker, E. P., *J. Math. Mech.* **17**, 59 (1967).

[11] Mermin, N. D., *Phys. Rev. Lett.* **65**, 3373 (1990).
Mermin, N. D., *Rev. Mod. Phys.* **65**, 803 (1993).

[12] Peres, A., *Phys. Lett.* **151**, 107 (1990).

[13] Hasegawa, Y. et al., *Nucl. Instr. & Meth. A* (doi:10.1016/j.nima.2010.06.234), 1-4 (2010).

[14] Oron, R., Shimshi, L., Blit, S., Davidson, N., Friesem, A. A. & Hasman, E., *Applied Optics* **41** (18), 3634-3637 (2002).

[15] Hasegawa, Y. et al, *Nature* **425**, 45-48 (2003).
Hasegawa, Y. et al, *Nucl.Instr. & Meth. A*, doi:10.1016/j.nima.2010.06.234 (2010).

[16] Pancharatnam, S., *Proc. Ind. Acad. Sc.* **44**, 247 (1956). Reprinted in *Collected Works of S. Pancharatnam*, ed. by G. W. Series (New York, OUP).
Ramaseshan, S. & Nitananda, R., *Current Sc* (India), **55**, 1225 (1986).

[17] Love, G. D., *Optics Comm.* **131**, 236-240 (1996).

[18] Sternberg, S., 'Lie Algebras', http://www.math.harvard.edu/∼shlomo/docs/lie_algebras.pdf (2004).